

CLAIM LANGUAGE:

1. (Currently amended) A recombinant antibody having a binding activity to 2,3,4,7,8-pentachlorodibenzofuran (2,3,4,7,8-PeCDF), which comprises ~~at least one polypeptide~~ a variable heavy chain or a variable light chain selected from the group consisting of:

(1) ~~a variable heavy chain a polypeptide constituting the H chain variable region of monoclonal antibody Dx3860 recognizing 2,3,4,7,8-PeCDF, and~~ having the amino acid sequence as shown in SEQ ID No. 5;

(2) ~~a variable light chain a polypeptide constituting the L chain variable region of said monoclonal antibody Dx3860, and~~ having the amino acid sequence as shown in SEQ ID No. 6;

(3) ~~a variable heavy chain a polypeptide constituting the H chain variable region of monoclonal antibody Dx3150 recognizing 2,3,4,7,8-PeCDF, and~~ having the amino acid sequence as shown in SEQ ID No. 7;

(4) ~~a variable light chain a polypeptide constituting the L chain variable region of said monoclonal antibody Dx3150, and~~ having the amino acid sequence as shown in SEQ ID No. 8; and

(5) a variable heavy chain polypeptides having amino acid sequences sequence as shown in any one of SEQ ID Nos. 64-67 ~~showing not less than 95% of homology to the amino acid sequences of the above polypeptides (1) (4),~~
and having a binding activity to 2,3,4,7,8-PeCDF; ~~and~~

~~—— (6) polypeptides representing fragments of the above polypeptides (1) (5), and having a binding activity to 2,3,4,7,8-PeCDF.~~

2. (Canceled)

3. (Currently amended) ~~The A recombinant antibody according to claim 1 which comprises:~~

~~a polypeptide constituting the H chain variable region of monoclonal antibody Dx3860 and having the a variable heavy chain region having the amino acid sequence as shown in SEQ ID No. 5, or a polypeptide having an amino acid sequence showing not less than 95% of homology to the amino acid sequence of the former polypeptide any one of amino acid sequences as shown in SEQ ID Nos. 64-67 and having a binding activity to 2,3,4,7,8-PeCDF; and~~

~~a variable light chain region polypeptide constituting the L chain variable region of monoclonal antibody Dx3860 and having the amino acid sequence as shown in SEQ ID No. 6, or a polypeptide having an amino acid sequence showing not less than 95% of homology to the amino acid sequence of the former polypeptide and having a binding activity to 2,3,4,7,8-PeCDF.~~

4. (Currently amended) ~~The A recombinant antibody according to claim 1 which comprises:~~

Serial Number: 10/550,951
Attorney Docket: SAWA3005/ESS

a variable heavy chain region~~polypeptide constituting the H chain variable region of monoclonal antibody Dx3150 and~~ having the amino acid sequence as shown in SEQ ID No. 7,~~or a polypeptide having an amino acid sequence showing not less than 95% of homology to the amino acid sequence of the former polypeptide and having a binding activity to 2,3,4,7,8-PeCDF; and~~

a ~~polypeptide constituting the L chain-variable~~ light chain ~~region of monoclonal antibody Dx3150 and~~ having the amino acid sequence as shown in SEQ ID No. 8,~~or a polypeptide having an amino acid sequence showing not less than 95% of homology to the amino acid sequence of the former polypeptide and having a binding activity to 2,3,4,7,8-PeCDF.~~

5. (Original) A DNA encoding the amino acid sequence of the recombinant antibody according to claim 1.
6. (Original) A cloning or expression vector comprising the DNA according to claim 5.
7. (Original) A transformant transformed with the cloning or expression vector according to claim 6.
8. (Currently amended) A process for preparing ~~the a~~ recombinant antibody ~~according to claim 1~~, which comprises cultivating the transformant transformed with an

Serial Number: 10/550,951
Attorney Docket: SAWA3005/ESS

expression vector according to claim 7 in a suitable medium, and recovering the recombinant antibody from the transformant or the medium.

9. (Currently amended) A method for immunologically capturing 2,3,4,7,8-PeCDF, which comprises ~~the use of~~ binding of the recombinant antibody according to claim 1, to 2,3,4,7,8-PeCDF.

10. (Currently amended) A method for immunologically determining 2,3,4,7,8-PeCDF in a sample, which comprises ~~the use of~~ binding of the recombinant antibody according to claim 1, to 2,3,4,7,8-PeCDF.